

**PRELIMINARY STUDY OF AGRICULTURE LAND USE AND
QUALITY OF WATER RESOURCES IN PALEODRAINAGE DELTA,
EASTERN-SOUTH AL- QATTARA DEPRESSION, EGYPT.**

Soliman.Y.R.A

Soils, Water and Environ. Res. Inst., Agric. Res. Center, Giza, Egypt

ABSTRACT

The Paleodrainage Delta soils are located between longitude 27° 00' to 31° 00' East and latitude 27° 00' to 30° 00' North, covering about 44000.0 km². This study aims to give a preliminary vision pertinent agriculture land evaluation and quality of water resources using GIS techniques, Land sat ETM images and digital elevation model (DEM).

Fifty five soil profiles were taken to represent the main physiographic units, and soil samples were performed for physical and chemical analyses. Also, water resource samples were collected from two wells and chemically analysed.

Data reveal that landscapes include five physiographic units within the contour lines of delta belts as follows:

- 1- Highest belt of the delta > 225 m a.s.l.
- 2- High belt of the delta 150 – 225 m a.s.l.
- 3- Medium high belt of the delta 75 – 150 m a.s.l.
- 4- Lower belt of the delta 0 – 75 m a.s.l.
- 5- Longitudinal dunes.

The studied soils could be classified into two Taxonomic orders; Entisols and Aridisols within 26 soil family levels. The surface features are mainly almost flat to gently undulating relief partly with rock outcrop. These soils are deep to moderately deep, sand to sandy loam texture locally with high gravel content or shallow depth. The lime and gypsum content were relatively high with more percentage in the medium-high and lower belt of the delta. Soil salinity varied widely from non-saline to extremely saline.

Data showed that high current suitability values were in very high belt of delta, whereas moderate potential suitability values were in some low belt of delta. While, low current and potential suitabilities were in sand dunes unit. The obtained classes could be changed to the better as potential suitability.

Data revealed that waters quality of the two wells, No. 1 and 2 are not suitable for drinking usage, however iron content is mostly above the critical limits for such use. On the other hand, waters of well No. 1 have marginally suitable class (C3-C2) for irrigation use and may cause more restrictions when

used for a long-term. While, waters of well No. 2 have highly suitable class (C1-C1) for crop irrigation without any restrictions, hence when used for a long-term.

Key Words : Paleodrainage Delta, GIS, Western Desert, Qattara Depression, Geomorphology.

Introduction

The identification of the land resources for agricultural development justifies the importance of producing a collective physiographic soil maps for building up database of land information system. The land resources may be divided into three groups: (1) very stable resources (climate, relief, geological formations), (2) moderately stable resources (soils and water, some artificial elements of the land), and (3) relatively unstable resources (vegetation and related biological features);(Vink,1975).

Physiographic approach can provide a good basis for explaining geomorphology through aerospace image interpretation (Goosen, 1967). The physiographic genesis was performed to find a land attribute illustration for a vast area, considering the parent rock and the inherited parent material, which can be traced by the paleo and recent drainage patterns as mediators between the highlands and lowlands. The availability of data sources in a digital form and increased capability of computers to handle large volumes of data have allowed to create attributes data of soils, as with soil survey. Spatial mapping usually involves the interpolation of point data across surfaces to depict condition at all positions on the land surface. Remote sensing should be used as an aid to distinguish landscape element. Qualitative models derived from modification of soil landscape models are efficient means of interpolation point data based on conceptual relationship between observation of the soil property or condition being mapped and easily observable landscape features (Peterson *et al.* 1996). Geographic Information System (GIS) is considered as organized collection of computer hardware, software, spatial and non-spatial data that can help users for the efficient capture, storage, update, manipulation, analysis and management of all geographically referenced information. Features on the map are linked to records in the database which contain a multitude of attributes and values. These components (digital map and database) serve as a storehouse of information. The map stores physical features and the database stores information about them. The result of having these two components linked is that both spatial data (map features) and attribute data can be queried and retrieved. This is important for planners, who rely on data about geographic space. Additional mathematical functions can allow statistical analysis of data, create new data and create predictive models.

The aims of this study were to using remote sensing data for identify the landforms, their soil attributes and quality of irrigation water as land resources, in

PRELIMINARY STUDY OF AGRICULTURE LAND USE42

a vast area that has important situation for development and introduce it to one of the agriculture extension project.

Location of study area

The study area are located between longitude 27° 00' to 31° 00' East and latitude 27° 00' to 30° 00' North, and occupies a vast region that covers about 10 482 000 fadden (44000.0 km²), from east El-Qattara depression to west of Gabel Qatrani and west of Dyrout aligning of El-Menia and El-Fayoum provinces (Fig.1).

Geology

Issawi and Mc Cauley (1992) attributed the origin of these Paleodrainage (undulating delta apex) to the downstream parts of the Qena system, which formed by northward lateral erosion in the early Oligocene as the Red Sea Mountains rise. The eroded materials removed from the Red Sea Mountains to the northwest of El Fayoum basin depositing a delta across El Fayoum in huge quantities of deep fossiliferous Oligocene Fluvial sands and gravel. **Albritton et al. (1990)** proposed that Qattara Depression originated as a stream valley that was dismembered by Karstic processes during the late Miocene epoch, and after word it was by Fluvial processes. A major stream issuing from the Gilf Kebir highlands in the south, showed northward may have done so by route through the Qattara to an exit near the head of the Ras Alam or Rum submarine canyon offshore near Alexandria **Map (1)**.

Issawi et al. (2001) stated that the part of the Paleodeltaic sediments represents the Miocene transgression reached to about 60 km south of Cairo, and its sediments make up the surface of the desert to the north of El Fayoum basin. The degradation first removed the Oligocene sands and conglomerates of Gabel Qatrani formation, and Eocene bedrock of the depression bottom. It was accomplished when the Mediterranean was desiccated the late Miocene and streams cut deep canyons toward the sea bed, and the final phase took place in the Quarternary.

Data in the Geological map of Egypt (1987) reveal that apex of delta belong to tertiary (Miocene, Oligocene and Eocene), Quaternary (Undivided Quarternary and sand dunes), while enter and pro delta belong to Eocene era only.

Climate

The climatic data of studied area (CLAC, 2010), show a very low annual precipitations or nearly absent (about 7.5 mm/y). The mean annual temperature is above 22.0°C with. Minimum temperature is recorded in January, while maximum temperature is recorded in August. According to USDA (2010) the studied area has hyperthermic temperature regime with aridic and torric soil moisture regime.

Water resources

Nubian Sandstone Aquifer System (NASA) occupying much areas of Egypt and extended to the border of westward direction into Libya, in the south and southwestward direction into Sudan and Chad, **Shata(1987)**. The water resources which recorded in the studied area are **demonated** by El-Moghra aquifer system which have an average thickness of 300m (200 million m³/y), and Salinity >3000 ppm, with associated aquifers such as fractured carbonate aquifers and Nubian Sandstone aquifer, **Allam et al. (2002)**.




Figure 1: Location map of the study area

Map (1) THE EGYPTIAN GEOLOGICAL SURVEY AND MINING AUTHORITY 1981.

To achieve the aforementioned target, analysis of landsat images covering the area under investigation was done to identify the different physiographic units. Also, fieldwork was performed to describe the soils developed on these physiographic units as well as to collect soil samples for laboratory analyses to recognize the soil properties, which were used as guidelines for soil classification, hence land suitability for irrigated agriculture.

GIS work

The Geographic information system (GIS) was used for analytical and data management features including data input for reprocessing and output for final maps and useful arrays of digitizing operations for provide vector or raster based modeling capabilities for overlay and buffer analysis creation. All vector layers were imported into GIS. GIS is used for design spatial modeling of the soil properties of study area .The average weight values were calculated for each property of each profile using all profile depth. Data acquired in year 2010 of landsat Enhanced Thematic Mapper (ETM) which have spatial resolution 28-30 m were used for **delineation** the physiographic units of the studied area. The physiographic analysis detailed by **Goosen (1967)**. Updating of the physiographic

PRELIMINARY STUDY OF AGRICULTURE LAND USE44

map was carried out using the American landsat 7 image (path 176 row 41, path 177 row 39, path 177 row 40, path 178 row 41, path 178 row 40, path 179 row 39, and path 179 row 40), Fig. 2, and map contour line Fig. 3. The hardware which were used to carrying out the work are: PC (I7), GPS, scanner, and printers, software have been using to construct the geographic data are, Arc Gis 9.3 and Erdas Imagine 9.1.

Fieldwork and sampling

The preliminary image-interpretation map was checked in the field by different ground observation points to confirm the boundaries of the physiographic units or to revise what were shifted.

The different physiographic units were represented through 55 soil profiles and 2 water samples. The soil profiles were dug to a depth 150 cm or to lithic contact (bedrock). Soil samples were collected to represent the different morphological variations throughout the soil profile layers and were described according to the nomenclature of **FAO (2006) and the Munsell Color (2010)**. The disturbed soil samples were air-dried, crushed, sieved through a 2 mm sieve, then, the obtained fine earth samples (less than 2 mm) were kept for laboratory analysis.

Longitudes and latitudes of the studied profiles as well as their elevations were defined in the field using GPS.

Laboratory analysis

Particle size distribution was determined using the International Pipette method (**Piper, 1950**) and sodium hexametaphosphate as dispersing agent as described by **Baruah and Barthakur (1997)**. Calcium carbonate content was measured using the Collin's Calcimeter method (**Wright, 1939**). Saturation soil paste extract and soil pH in the soil water suspension of 1:2.5 were determined according to the methods described by **Jackson (1973)**.

Gypsum content, and the exchangeable sodium percentage were determined according to the methods described by **Richards (1954)**. Soils under study were classified into taxonomic units according to Soil Taxonomy System (**USDA, 1975**) and using the keys to Soil Taxonomy (**USDA, 2010**). Soils under investigation were evaluated using the parametric system for irrigated agriculture land evaluation namely "land Capability Classification" undertaken by **Sys and Verheye (1978)**, water resource samples were collected from two wells and chemically analysed. Well No.1 is located in 29° 09' 37" E & 28° 43' 23" N, having + 119 a.s.l. (above sea level), Well No.2 is located in 29° 26' 36" E & 28° 33' 47" N, having + 101 a.s.l. Water quality and its suitability have been compared with the standard guide line values of both drinking and irrigation water uses, issued by **Richards (1954), El-Ghandour et al. (1983), Ayers and Westcot**

(1985), Egyptian Ministry of Health (1995), Ministry of Environment (1999) and WHO (2008).




Figure 2: Landsat TM 7 image of the studied




Figure 3: shows contour lines in the study area

Results and Discussion

Data cited from **Issawi and McCauley (1992)**, and **Albritton et al. (1990)**, which illustrated previously in introduction, reveal to the studied area have different parent materials and multi origin.

On the other hand, the geological map of Egypt 1981(map1), cleared that studied area belong to five geological eras, while enter and prodelta belong to one era (Eocene).

Afify (2009) concluded that the studied area is a Paleodrainage delta, and the delta consist of three units, i.e. delta apex, inter delta and prodelta but have the same parent material and precipitate in the same era.

Delta apex, enter delta and prodelta have a descending elevation, respectively while, delta recorded by 34m to 311m a.s.l. Elevation model program appear that inter delta is higher than delta apex in most area, return to Figure (3). We can concluded that the studied area have not the pattern of the delta but we can describe it as a false delta, because it have the delta form only.

Tracing the landscape genesis for very **vast** area, such as the studied one, begin from Gable Qatrani Formation which its elevation tends to decrease gently to reach sea

level at El-Quattara depression in its west side, which intimate that the delineated main physiographic units are :

1- Highest belt of the delta (> 225 m a.s.l.)

PRELIMINARY STUDY OF AGRICULTURE LAND USE46

- 2- High belt of the delta (150 – 225 m a.s.l.)
 - 3- Medium-high belt of the delta (75 – 150 m a.s.l.)
 - 4- Lower belt of the delta (0 – 75 m a.s.l.)
 - 5- Longitudinal dunes
- a.s.l. = (above sea level)

The main physiographic units are illustrated in map (2), the morphological description of the studied soil profiles are summarized in Table (1), while physical and chemical attributes are recorded in Table (2). The delineated physiographic units are described as follows:-

1- Highest belt of the delta (> 225 m a.s.l.)

Soils of this unit are located in the central part of the study area surrounded with soils of high belt of delta and mountains. The surface features are characterized by almost flat to gently undulating relief and covered mainly by dark coloured coarse fragments as gravel. Most of the examined soil profiles are deep to moderately deep sandy to sandy loam texture mainly in subsurface reddish colour (7.5YR and 5YR). The clay content ranges of the most between 3 and 15% and the gravel content vary widely from nil to more than 50%. Soil structure is massive in slightly hard to very hard consistence by depth. The lime and gypsum content range from 2 to about 20% and from <1 to about 15%, respectively, that were observed in the field as few to common secondary accumulation. The soil reaction (pH) range from 7.6 to 8.2, that indicate moderate alkaline while soil salinity as determined by the Electrical conductivity, values (ECe) range from very slightly saline (2 to 4 dS/m) to strongly saline (16 to 32 dS/m)

By using the **Soil Taxonomy Key manual (USDA, 2010)**, the examined profiles in these soils were classified to the family level and the following taxonomic units were indentified as an associations kind:

-Weakly developed soils (Entisols order):

- Sandy soils: *Typic Torriorthents*, sandy, mixed, hyperthermic.
- Gravelly loam soils: *Typic Torriorthents*, loamy-skeletal, mixed, hyperthermic.

- Soils developed under arid conditions (Aridisols order):

- Lime accumulated /gravelly loam soils:
:*Typic Haplocalcids*, loamy-skeletal, mixed, hyperthermic.
- Calcigypsy sandy or gravelly sand soils:
:*Typic Calcigypsids*, sandy, mixed, hyperthermic.
- :*Typic Calcigypsids*, sandy-skeletal, mixed, hyperthermic.

And the following taxonomic units as inclusions:

- Lime accumulated sandy or gravelly sand soils:
:*Typic Haplocalcids*, sandy, mixed, hyperthermic.
- :*Typic Haplocalcids*, sandy-skeletal, mixed, hyperthermic.
- Gypsiferous gravelly loamy soils:
:*Typic Haplogypsids*, loamy skeletal, mixed, hyperthermic.

Table (1)

PRELIMINARY STUDY OF AGRICULTURE LAND USE48

Table (1) Cont.

Table (1) Cont.

PRELIMINARY STUDY OF AGRICULTURE LAND USE50

Table (1) Cont.

Table (1) Cont.

PRELIMINARY STUDY OF AGRICULTURE LAND USE52

Table (1) Cont.

Physiographic units	Profile No.	Depth (cm)	Gravel %	Color		Structure	Consistency (Dry)	Effervescence	Secondary Formation %		Boundary
				dry	moist				CaSO4. 2H2O	CaCO3 %	
Lower belt of the delta 0 – 75 m a.s.l	43	0 - 20	0.0	10YR7/6	10YR6/6	m	s	mo	Very few	few	as
		20-50	0.0	10YR7/3	10YR6/4	m	s	sl	many	few	
	44	0 - 20	0.0	10YR7/6	10YR6/8	m	l	st	Very few	few	as
		20-45	0.0	10YR5/4	10YR5/8	m	h	mo	many	few	
	45	0 - 25	5.0	10YR7/6	10YR6/6	m	s	st	Very few	few	aw
		25-70	0.0	10YR7/4	10YR6/4	m	h	sl	many	few	
	46	0 - 25	15.0	10YR6/4	10YR5/6	m	sh	st	Very few	common	cs
		25-65	10.0	7.5YR7/6	7.5YR8/6	m	sh	st	many	common	cw
		65-130	0.0	7.5YR7/6	7.5YR6/6	m	v.h	st	many	many	
	47	0 - 20	10.0	10YR6/4	10YR5/6	m	s	st	few	many	cw
		20-70	25.0	10YR6/4	10YR5/6	m	s	mo	many	many	gw
		70-130	5.0	7.5YR7/6	7.5YR6/4	m	v.h	sl	many	many	

Table (1) Cont.

Physiographic units	Profile No.	Depth (cm)	Gravel %	Color		Structure	Consistency	Effervescence	Secondary formation		Boundary
				dry	moist				CaSO4 . 2H2O	CaCO3 %	
Longitudinal dunes.	35	0 - 150	0.0	10YR ^v /6	10YR6/6	sg	lo	sl	-	-	-
	48	0 - 150	0.0	10YR ^v /6	10YR6/6	sg	lo	sl	-	-	-
	55	0 - 150	0.0	10YR ^v /6	10YR6/6	sg	lo	sl	-	-	-

Soil structure: gr=granular(single grain), bw: blocky weak grain , bm: blocky medium, pm: platy medium. .
Consistency: v.h=very hard,h=hard, f=friable, lo=loose, so=soft, vs=very sticky, ss=slightly sticky, vp=very plastic and sp=slightly plastic.
Effervescence: st: strong, mo:moderate, sl: slightly effervescence.
Consistency: vh= very hard, h=hard, sh= slightly hard, so=soft, lo=loose.
Lower boundary: cs=clear smooth, cs: clear smooth, cw: clear wavy, aw : abrupt wavy, , as : abrupt smooth, and gw: gradual Wavy .

PRELIMINARY STUDY OF AGRICULTURE LAND USE54

Table (2)

Table (2) Cont.

PRELIMINARY STUDY OF AGRICULTURE LAND USE56

Table (2) Cont.

Table (2) Cont.

Physiographic units	Profile No.	Depth (cm)	Grain size distribution %			Modified Texture Class	pH (1:2.5)	EC (dS/m)		CaSO ₄ . 2H ₂ O %	CaCO ₃ %
			Sand %	Slit %	Clay %			Soil past	ESP		
Lower belt of the delta 0 – 75 m a.s.l	43	0 - 20	89.47	6.21	4.32	S	7.71	30.67	24.90	0.12	3.49
		20-50	46.32	24.34	29.34	SCL	7.98	48.54	54.94	15.25	2.73
	44	0 - 20	90.51	6.45	3.04	S	8.69	81.11	33.54	0.21	1.62
		20-80	46.90	19.89	33.21	SCL	7.83	167.93	50.93	33.53	0.85
	45	0 - 25	88.71	7.08	4.21	S	7.9	8.78	6.26	0.19	5.79
		25-70	50.69	19.67	29.64	SCL	7.89	9.29	14.07	16.16	2.98
	46	0 - 25	79.49	13.64	6.87	LS	7.55	46.76	42.13	0.54	10.00
		25-65	79.93	13.02	7.05	LS	7.82	37.96	33.24	16.44	13.35
		65-130	25.85	29.37	44.78	C	7.27	39.32	35.28	17.51	19.61
	47	0 - 20	78.12	14.87	7.01	LS	7.89	54.44	39.97	2.90	29.29
		20-70	79.64	13.08	7.28	LS	7.73	48.98	47.89	24.85	28.02
		70-130	22.56	32.67	44.77	C	7.39	43.76	31.72	19.53	29.80

Table (2) Cont.

Physiographic units	Profile No.	Depth (cm)	Grain size distribution %			Modified Texture Class	pH (1:2.5)	EC (dS/m)		CaSO ₄ . 2H ₂ O %	CaCO ₃ %
			Sand %	Slit %	Clay %			Soil past	ESP		
Longitudinal dunes.	35	0 - 150	92.68	4.68	2.64	S	8.2	2.23	4.68	0.00	0.00
	48	0 - 150	89.03	7.64	3.33	S	8.15	1.08	1.33	0.43	0.00
	55	0 - 150	91.96	6.03	2.01	S	8.2	2.23	4.37	0.15	0.52

Fine earth: S=Sand, LS=Loamy sand, SL=Sandy loam, SCL=Sandy clay loam, CL=Clay loam and C=Clay.

2- High belt of the delta (150 – 225 m a.s.l)

Soils of this unit are located in the central part of the study area surrounded with soils of highest belt of delta and bordered from north and south with medium-high belt of the delta. The surface features and soil attributes are almost similar to that of highest belt unit but differ in the range of some soil characteristics. The topography is relatively less undulating and also for surface dark coloured gravel. The examined profiles indicate more coarser texture in most locations but in less gravel content and locally with shallow soil depth over rock. Both lime and

PRELIMINARY STUDY OF AGRICULTURE LAND USE58

gypsum seem to be in relatively lower content, but soil salinity ranges mainly from moderately (8 to 16 dS/m) to very strongly saline (32 to 64 dS/m).

According to the characteristics of the examined soil profiles, the following association taxonomic units were identified:

-Weakly developed soils (Entisols order):

- Sandy or Gravelly sand soils: *Typic Torriorthents*, sandy, mixed, hyperthermic.

:*Typic Torriorthents*, sandy-skeletal, mixed, hyperthermic.

-Loamy or Gravelly loam soils: *Typic Torriorthents*, coarse-loamy, mixed, hyperthermic.

:*Typic Torriorthents*, loamy -skeletal, mixed, hyperthermic.

- Soils developed under arid conditions (Aridisols order):

- Lime accumulated sandy soils:

:*Typic Haplocalcids*, sandy, mixed, hyperthermic.

- Gypsiferous sandy or gravelly sand soils:

:*Typic Haplogypsids*, sandy, mixed, hyperthermic.

:*Typic Haplogypsids*, sandy -skeletal, mixed, hyperthermic.

:*Leptic Haplogypsids*, sandy, mixed, hyperthermic.


And the following taxonomic units as inclusions:

- Shallow sandy soils: :*Lithic Torriorthents*, sandy, mixed, hyperthermic.

- Shallow Gypsiferous gravelly sandy soils:*Lithic Haplogypsids*, sandy-skeletal, mixed, hyperthermic.

- Calcigypsy gravelly sand soils:

:Typic Calcigypsids, sandy-skeletal, mixed, hyperthermic.



Map. (2): Physiographic units map of the studied area- Eastern-South Desert Outskirt of Al-Qattara depression, Egypt

3- Medium- high belt of the delta (75 – 150 m a.s.l.)

Soils of this unit are surrounding the high belt unit and bordered from north with lower belt unit. The surface features are characterized by almost flat relief covered by common relatively light coloured gravel. Also, the surface is mainly

PRELIMINARY STUDY OF AGRICULTURE LAND USE60

dissected by many shallow gullies and the scattered rock outcrops were observed. The depth of the examined soil profiles vary from deep to shallow over rock or indurated layer. The soil texture is dominated by sandy to sandy loam in 10YR to 7.5YR hue colour and mixed with high gravel content in some profiles. The relatively high lime and gypsum content were indicated in the most of soil layers that observed in the field as common to many secondary accumulation with indurated gypsum in some subsoil layers. The soils are moderately alkaline and salt content varied widely from slightly to very strongly saline. Accordingly, the following taxonomic unit could be identified the studied soil profiles:

- Calcigypsy gravelly loam soils:

Typic Calcigypsids, sandy-skeletal, gypsic, hyperthermic.

- Calcigypsy sandy soils: *Typic Calcigypsids*, sandy, mixed, hyperthermic.

- Shallow Gypsiferous soils:

- *Calcic Petrogypsids*, fine loamy, gypsic, and hyperthermic.

- *Lithic Calcigypsids*, sandy, carbonatic, hyperthermic.

- Lime accumulated gravelly soils:

- *Typic Haplocalcids*, loamy-skeletal, mixed, hyperthermic.

- *Typic Haplocalcids*, sandy-skeletal, mixed, hyperthermic.

-And the following taxonomic units as inclusions:

- Weakly developed soils: *Typic Torriorthents*, sandy, mixed, hyperthermic.

- *Typic Torriorthents*, fine loamy, mixed, hyperthermic.

-Gypsiferous sandy soils:

- *Typic Haplogypsids*, sandy skeletal, gypsic, hyperthermic.

4- Lower belt of the delta (0 – 75 m a.s.l)

This unit is found in the northern portion of the study area. The surface features are dominated by almost flat to gently undulating relief covered with few to common dark coloured gravel. The soil profiles are characterized by deep to moderately deep loamy or clayey texture mainly in 10YR hue colour, with coarse texture in the surface layers. Soil structure is massive mainly in hard and very hard consistence. The subsurface layers indicate high gypsum content with lime varied widely from 3 to 30%. The soils are very strongly to extremely saline with moderately soil reaction. The soil profiles indicate the following taxonomic units:

- Gypsiferous loamy soils: *Typic Haplogypsids*, fine loamy, mixed, hyperthermic.

- Calcigypsy soils: *Typic Calcigypsids*, sandy over clayey, gypsic, hyperthermic .

- Salt accumulated soils: *Gypsic Haplosalids*, fine loamy, gypsic, hyperthermic .

5- Longitudinal dunes

This unit are located over all previous units. It formed by the action of windblown sand as high longitudinal dunes that extend from the west-north to east

south direction. It characterized by homogeneous yellow loose sand almost free from lime and gypsum that classified as : -*Typic Torripsammets*, siliceous, hyperthermic.

Land suitability Evaluation:

Land evaluation was calculated based on the concepts of Sys & Verheye (1978), data in Table (3) show that current and potential suitability classes vary between Non suitable and highly suitable with dominance low suitability levels, the soil Texture including gravel is the main factor for decreasing the suitability class, followed by soil salinity and alkalinity while lime and gypsum content as well as soil depth indicate different severity degree . The higher suitability values were in highest belt of delta in the current state and in lower belt of delta in the potential state because of its high salinity levels which can be fixed by using the optimum cultivation methods, while the lowest values of current and potential suitabilities were in sand dunes unit. The major land improvement accelerates the utilization of the soil under consideration as follow:

- a-** Using the organic matter as the soil conditioner and fertilizers.
- b-** Application system of irrigation control of water percolation and wetting the root zone only such as drip irrigation, for sprinkler irrigation systems decreases the importance of soil texture as a main factor of land evaluation.
- c-** Selection more suitable crops which are in agreement with soil components.

Water quality and its Suitability:

Data in Table (4) showed that the TDS and major ions content of the sampled wells (No.1 and No.2) are below the critical limits for drinking water guide line values, but the trace element content of ions is mostly above the critical limits of such use according to **Ministry of Health (1995)** and **WHO (2008)**. Therefore, water resources of the two wells are not suitable for drinking usages.

According to **Richards (1954)** and **El-Ghandour et al.(1983)**, the sampled water corresponding to their TDS, EC and SAR values are characterized by high salinity levels with medium alkalinity hazard (C₃-S₂); having marginally suitable class for irrigation use, (well No.1). Therefore, it may cause more restrictions when used in irrigation for a long-term. On the other hand, the sampled waters of well No.2 are characterized by low salinity levels with low alkalinity hazard (C₁-C₁), having highly suitable class for crop irrigation, hence when used for long-term.

According to **Ayers Westcot (1985)**, water quality of well No.1 appears increasing problem level of salinity (1.2 dS/m) effect on grown crops, while its effect on soil permeability is no problem while values of adj SAR>9.00 reveal to sever problem of permeability and sever problem of sodium toxicity, increasing problem for chloride ions specially Foliar absorption from leaves wetted by sprinkler applied water. No problem level due to boron contents (0.193 ppm).

Table (3)

Table (3): Cont.

PRELIMINARY STUDY OF AGRICULTURE LAND USE64

Table (3): Cont.

Table (3): Cont.

PRELIMINARY STUDY OF AGRICULTURE LAND USE66

Total contents of nitrogen (NO_3^- & NH_4^+) < 5 ppm reveal to no problem. Bicarbonate contents (4.85 me/L) may cause problems while deposits on fruit or leaves due to sprinkler irrigation. Values of pH in normal range, while, pHc values <8.4 indicate tendency to precipitate lime from waters applied.

Value of residual sodium carbonate >2.5 is considered not suitable for irrigation according to **Van Hoorn (1971)** where a high content of bicarbonate in irrigation water may lead to precipitation calcium and magnesium in the soil and thus to a relative increase of the sodium concentration.

Contents of soluble trace elements are as follow Iron 1.17 ppm Manganese 0.086 ppm, Zinc 0.005 ppm, and Copper nil. However the soil type should be taken in the consideration, whereas, most of soil under consideration sandy or gypsiferous soils.

The sampled waters of well No. 2 show that salinity and adj. SAR values appear no problem, for soil permeability, or specific ion toxicity for plant roots and foliar absorption.

Total contents of Nitrogen (NO_3^- & NH_4^+) have no problem (1.7 ppm), pH value in a normal range, while Bicarbonate contents reveal to increasing problem level. Residual sodium carbonate appears no-sodium hazard.

Soluble trace elements contents are present as follows: Fe 0.673 ppm, Mn 0.115 ppm, Zn 0.001 ppm and Cu 0.001 ppm. Therefore the second water well considers a good quality and suitability, when used for irrigation use for a long-term.

Table (4a): Some chemical analyses of the studied water resources.

Well No.	pH	Soluble cations				Soluble anions				Boron	Zinc	Copper	Manganese	Iron	N	N	TSD (ppm)	pHc	SAR	adj. SAR	
		(meq/L)				(meq/L)															
		EC (dS m ⁻¹)	Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	CO ₃ ⁼	HCO ₃ ⁻	Cl ⁻												SO ₄ ⁼
1	7.5	1.20	0.74	1.01	9.65	0.62	0.0	4.85	6.01	1.20	0.193	0.005	0.00	0.086	1.170	0.95	1.89	768.0	7.58	3.06	18.782
2	7.4	0.39	0.84	0.69	2.34	0.13	0.0	2.21	1.02	0.77	0.032	0.001	0.001	0.115	0.673	1.70	0.0	250.0	8	0.68	3.744
Well location		Salinity classification				Permeability Classification				pH		Specific ion toxicity									
1		(Ip)				EC ----- (Ip) Adj SAR - (Sp)				(Nr)		(Np) for Na, Cl, Zn, M, Fe, Cu and B									
2		(Np)				EC ----- (Np) Adj SAR - (Np)				(Nr)		(Np) for Na, Cl, Zn, Mn, Fe, Cu and B									

Np Np =No problem ,IP= Increasing problem,SP= Sever problem, RSC=Residual sodium carbonate, SAR= Sodium adsorption ratio, adj. SAR= adjusted sodium adsorption ratio, Nr= Normal range

REFERENCES

- Affy,A.A. (2009).** Land resources evaluation of of the Paleodrainage delta in Western desert of Egypt using remote sensing data.Egypt.J.Remote Sensing & Space sci.V12. pp 3-26.
- Albritton, C. C.;Brooks, J.E.I.;Issawi,B.and Swdan, A. (1990).** Origin of the Qattara Depression, Egypt. Geological Society of America Bulletin 102 (7), pp. 952-960.
- Allam A., E. Saaf and M. Dawoud (2002).** Desalination of brackish groundwater in Egypt. Desalination, Vol. 152: 19-26.
- Ayers,R.and D.W.Westcot (1985).** Water quality for agriculture.(FAO Irrigation and Drainage Paper 29,Rev.1) Rome:FAO,164 pp.
- Baruah, T.C. and H.P. Barthakur (1997).** A Text Book of Soil Analysis. Vikas Publishing Housing, PVT LTD, New Delhi, pp. 35-67.
- CLAC, (2010).** Central Laboratory for Agriculture of Climate. Web site www.clac.edu.eg.
- Egyptian Ministry of Health (1995).** Standards and specifications of water quality for drinking and domestic uses.

PRELIMINARY STUDY OF AGRICULTURE LAND USE68

- El-Gandour, M.F., K. Hefny, J.B. Khalil and S.A. Atta(1983).** Suitability of groundwater in the Nile Delta region(Egypt) for irrigation. Middle East water & Sewage, 7(2), Fuel & Metallurgical Jout, Ltd. U.K.
- FAO (1994):** "Guideline For Soil Profile Description". FAO ISRIC Publication, Rome.
- FAO (2006).** Guidelines for soil profile description. FAO, ISRIC Publication, Rome.
- Goosen, A.A.I. (1967):** "Aerial photo-interpretation in soil survey". Soil Bull. No. 6, FAO, Rome, Italy.
- Issawi,B. and Mc Cauley,J.F. (1992).** The Cenozoic rivers of Egypt : The Nile problem : in Friedman, R. and Adams ,B.(eds),The followers of Horus : Studies in memory of M.A.Hoffman:Egption Studies Association Publication No.2,Oxbow Monograph 20, Oxbow Books, Oxford, England, pp. 121-138.
- Jackson, M.L. (1973).** Soil Chemical Analysis. Prentice India Private LTD, New Delhi, Indian.
- Ministry Of Industry and Mineral Resources 1981."** The Egyptian Geological Survey and Mining Authority".
- Ministry of the Environment (1999).**Guidelines for irrigation water quality. Human Resource Development and Employment, Department of Environment.
- Munsell Soil Color Charts (2010).** Edition Munsell Color, Macbeth Division of Kollmorgen Corp., 2441 North Calvert Street, Baltimore, Maryland, USA.
- Peterson, G.W., F.F. Connors, D.A. Miller and R.L. Gardner, 1996:** Aircraft and satellite remote sensing of desert soils and landscapes. Remote Sensing Environ. No, 23: 253-271.
- Piper, G.S. (1950).** Soil and Plant Analysis. A Monograph from the Wails Agric. Research Inst., University of Adelaide Australia.
- Richards, L.A. Ed. (1954).** Diagnosis and Improvement of Saline and Alkali Soils. U.S.D.A. Hand Book No. 60.
- Said, R. (1990):** "The Geology of Egypt". Published for the Egyptian Central Petroleum Corporation, Conoco Hurchada, Inc. by Balkema, A.A., Rotterdam, the Netherlands.
- Shata,A.(1987).** "Management problems of the Major Regional Aquifer in N.E. Africa." UN Tech workshop. Khartoum.

- Sys, C. and W. Verheye (1978): An attempt to the evaluation of physical land characteristics for irrigation according to the FAO framework for land evaluation. Int. Train Cent. Post Grad. Soil Sci. Ghent, Belgium pp 66-78.
- The Egyptian General Petroleum Corporation(1987).Geological map of Egypt (Scale 1: 500000). Sheet No. NG36 SW Luxor.
- U.S.D.A. (1975). Soil Taxonomy Basic System of Soil Classification. U.S. Dept., Agric. Handbook No. 436, USDA. Washington, D.C. 20402.
- U.S.D.A. (2010):Keys of Soil Taxonomy , 11 Edition , By Soil Survey Staff.
- Vink, A.P.A. (1975): Aerial photographs and the soil sciences. UNESCO, Proc. Toulouse Conf., Canada.
- Wright, C.H. (1939): Soil Analysis. Thomas Murby & Co., London.
- World Health Organization(WHO)(2008).Guidelines for drinking water quality (electronic resource): incorporating first addendum. Vol.1, Recommendations 3rd ed. Geneva, Switzerland, 595 pp.

دراسة أولية لإستخدام الأرض للزراعة ونوعية مصادر المياه في دلتا الصرف القديم
جنوب شرق منخفض القطارة - مصر

ياسر ربيع أمين سليمان

معهد بحوث الأراضي والمياه والبيئة – مركز البحوث الزراعية – جيزة - مصر

تمتد اراضي دلتا الصرف القديم من شرق منخفض القطارة الي غرب جبل قطراني والي الغرب بمحاذاة محافظتي الفيوم والمنيا حتي ديروط ، حيث تشغل منطقة الدراسة مساحة شاسعة تغطي حوالي ١٠٤٨٢٠٠٠ فدان تقريبا (٤٤٠٠٠ km²) . وتهدف الدراسة الي إعطاء رؤية أولية لتقييم الأرض الزراعية وتحديد نوعية مصادر المياه باستخدام تقنيات نظم المعلومات الجغرافية (GIS) و تكتيك التفسير المرئي لصور الأقمار الفضائية (7 Enhanced Thematic Mapper)، ونموذج الارتفاعات الرقمية (DEM) في انتاج خريطة عامة للاشكال الارضية كوحداث فيزيوجرافية تم تسميتها كالآتي:

١- حزام الدلتا المرتفع جدا (اكبر من ٢٢٥ م فوق سطح البحر).

٢- حزام الدلتا المرتفع (من ١٥٠ - ٢٢٥ م فوق سطح البحر).

٣- حزام الدلتا متوسط الارتفاع (من ٧٥ - ١٥٠ م فوق سطح البحر).

٤- حزام الدلتا المنخفضة (من ٠ - ٧٥ م فوق سطح البحر).

٥- الكثبان الرملية الممتدة.

وقد تم تحديد ودراسة اكثر من مائة وخمسون نقطة ملاحظة أرضية لوضع دليل ملامح الوحدات الخريطية (الوحدات الفيزوجرافية الرئيسية)، ثم حددت الاختلافات بين هذه الوحدات من خلال دراسة خمسة وخمسون قطاعا أرضيا تم وصفها مورفولوجيا وأخذت منها عينات تربة للتحليلات المعملية.

وتشير النتائج ان تحديد مظاهر السطح الذي يمثل الوحدات الفيزوجرافية كان متمشيا مع خطوط الكنتور المختلفة الارتفاعات عن سطح البحر لتمثل احزمة الدلتا القديمة.

PRELIMINARY STUDY OF AGRICULTURE LAND USE70

وقد امكن تقسيم الاراضي المدروسة الي رتبتي (الاراضي التي تطورت تحت ظروف الجفاف) Aridisols، (الاراضي حديثة التكوين) Entisols متضمنة ستة وعشرون وحدة تقسيمية حتي مستوي العائلة – ويتضح ان مظاهر السطح بها ان مظاهر السطح بها تميل الي طبوغرافية شبة مستوية الي خفيفة التموج يتخللها جزئيا بروزات صخرية في بعض الوحدات وقطاعات التربة اتصفت غالبا بقوام رملي الي طمي رملي عميقة الي متوسط العمق بعضها ذات محتوى حصوي مرتفع او ضحلة العمق . والتربة غالبا تحتوي علي نسب مرتفعة نسبيا من الجير والجبس وتكون اكثر وضوحا في المناطق المتوسطة الارتفاع والاكثر انخفاضاً من الدلتا ، كما تباينت ملوحة التربة كثيرا من الغير ملحية الي شديدة الملوحة. وتوضح النتائج ان قيم درجات صلاحية الارض الحالية للزراعة يمثلها حزام الدلتا المرتفعة جدا والمرتفعة، في حين تمثل اعلي قيم متوسطة الصلاحية للأراضي المتوقعة لبعض أراضي حزام الدلتا المنخفضة – بينما اقل قيم الصلاحية الحالية والمتوقعة كانت لوحدة الكتبان الرملية، ويمكن القوا بأن درجات الصلاحية الحالية يمكن الوصول بها الي درجة الصلاحية الافضل لتمثل الحالة المتوقعة.

وقد اثبتت النتائج ان نوعية مياه الآبار رقم (1) و (2) غير صالحة للشرب وذلك لإرتفاع المحتوي من عنصر الحديد بها عن الحد المسموح لأستخدامة ، ومن ناحية اخري فان درجة نوعية مياه البئر رقم 1 تكون حدية الصلاحية للري وقد تسبب اضرارا بالغة عند استخدامها للري لفترات طويلة. بينما تمثل درجة نوعية مياه البئر رقم (2) صلاحية عالية لري المحاصيل ولا تسبب اضرار متوقعة حتي مع الاستخدام لفترات طويلة، وذلك باتباع النظام الكمي الرياضى الموضوع بمعرفة (Sys and Verheye (1978).