

RESPONSE OF TWO MAIZE (*Zea mays* L.) GENOTYPES FOR SOME WATER STRESS AND NITROGEN FERTILIZATION TREATMENTS

F.S. Abd El-Samei, M.D.H. Dewdar and S.K.A. Ismail
Argon. Dept. Fac. Agric., Fayoum Univ.

ABSTRACT

Two field experiments were conducted during 2006 and 2007 successive seasons at the experimental Farm of Fayoum Fac. of Agric., to quantify the single and combined effects of different irrigation intervals (10, 15 and 20 days) and rates of nitrogen fertilization (100, 120 and 140 kg N/fed.) on growth, yield and yield components of two maize genotypes, i.e. (single-cross 10) and (three-way cross 310). The results indicated that S.C.10 gave the highest values of most studied characters, especially plant height, 100-grain weight and grain yield (kg/fed.) where the values were 245 (cm), 35.97 (g) and 2475.37 (kg), respectively.

Application of 10 days (I_1) significantly increased plant height, ear length, ear diameter, 100-grain weight and grain yield/feddan. Using the rate of 140 kg N /fed. (N_3) significantly increased plant height, leaf area, number of rows /ear, number of grain, 100-grain weight and grain yield/feddan by values of 262 (cm), 2678 (cm²), 13.30 (row), 44.64 (grain), 37.53 (g) and 2565.17 (kg) surpassing 100 (N_1) and 120 kg N/fed. (N_2). The results also indicated that planting S.C.10 and T.W.C. 310 under 10 day (I_1) gave the lowest number of days to 50% tasseling and silking traits.

Key words: Maize hybrids, irrigation intervals, nitrogen fertilization

INTRODUCTION

Maize is one of the most important cereal crops used for human consumption and many other purposes such as animal feeding and various industrial intentions. Recently, the national policy is to mix wheat flour (80 %) with maize flour (20 %) in making bread all over the country in order to reduce wheat grain imports. Therefore, in recent years, great efforts have been directed to increase maize production by planting high yielding genotypes under improved cultural practices. Such efforts reflected in a visible national improvement in maize productivity. However, at Fayoum Governorate, this productivity is still relatively low because of some farmers are frequently cultivate maize using their own types with unsuitable dose of nitrogen fertilization especially in the areas suffering from irrigation water shortage. Factors such as varieties / hybrids, irrigation intervals and rates of nitrogen fertilization plays a great role in maize production. It is very important to determine the best irrigation interval, dose of nitrogen fertilization and the high yielding potential hybrids for maximizing of grain yield.

The S.C.10 surpassed T.W.C.310 in plant height and leaf area (**Atta-Allah, 1996 and Sharaan et al., 2002 a**). Short irrigation interval increased plant height and leaf area/plant (**El-Ganayni, 2000**) while missing one irrigation significantly reduced plant height and area of leaves (**Abo-El-Kheir and Mekki, 2007**). Increasing irrigation intervals caused a reduction in no. of days to 50% tasseling and silking (**Ashoub et al, 1996 and Sharaan et al., 2002a**), and grain yield (**El-**

Ganayni, 2000). Single- cross-10 had superiority in grain yield over the three way cross 310 (**El- Sheikh, 1999 and Sharaan et al, 2002 b**). Growth and grain yield of different maize varieties increased with increasing in N levels (**Sharar et al, 2003**).

The present investigation was designed to study the responses of two maize hybrids to different irrigation intervals and nitrogen doses. In addition to determine the best combination among the studied of factors to produce the maximum yield under Fayoum condition.

MATERIALS AND METHODS

Two field experiments were carried out at the farm of Fayoum Fac. of Agric., during 2006 and 2007 seasons to study the responses of two maize genotypes to different irrigation intervals and nitrogen doses. A split-split-plot design with four replications was used. The soil was clayey with pH value of 7.5, organic mater of 1.63%, available nitrogen, phosphorus and potassium (30.22, 6.90 and 591 ppm), respectively. The experimental treatments were as follows:

Genotypes (main-plots):

G₁:Single cross 10 (S.C.10).

G₂:Three way cross 310 (T.W.C.310).

Irrigation intervals (sub-plots):

I₁: irrigation every 10 days.

I₂: irrigation every 15 days.

I₃: irrigation every 20 days.

Nitrogen fertilization (sub-sub-plot):

N₁:100 kg N/feddan.

N₂:120 kg N/feddan.

N₃:140 kg N/feddan.

The plot size was 21 m² (6.0 x 3.5 m) containing five ridges of 6.0 m length and 0.70 m width. Plots were isolated by border from all sides to avoid the effect of lateral movement of irrigation water. Calcium super phosphate (15.5 % P₂O₅) was added during field preparation at the rate of 100 kg / feddan. Nitrogen fertilization (Ammonium nitrate 33.5% N) was applied in two equal doses (before the first and second irrigation). Irrigation interval treatments were practiced after the second irrigation. Seeding rate was 15.0 kg / feddan and grains were planted on May 28th and June 4th in first and second seasons, respectively. The other recommended cultural practices for growing maize were adopted from planting till harvesting. The studied characters were plant height (cm), leaf area /plant (cm²), number of days to 50% tasselling, number of days to 50% silking, ear diameter (cm), ear length (cm), number of rows/ear, number of grains/ row, 100 grain weight (g) and grain yield (kg/fed.), the later trait was adjusted to 15.5% moisture content.

The obtained data were subjected to analysis of variance and the means were compared using LSD (0.05), according to **Gomez and Gomez (1984)**.

RESULTS AND DISCUSSION

1-Growth measurements

Data presented in Table (1) indicate that plant height and leaf area per plant were significantly affected by genotypes, irrigation intervals, nitrogen fertilization. While their interaction effects on both traits were insignificant indicating that they acted independently. Maize genotypes showed significant differences in plant height and leaf area per plant but each genotype had superior trait, where G₁ (S.C.10) had the tallest plant (245 cm) compared with G₂, while G₂ (T.W.C.310) had the greatest area of leaves per plant (2813 cm²) compared to G₁. These result may be due to the genetical differences between the two genotypes and are in the same trend with those obtained by Sharar *et al* (2003) and Amanullah *et al* (2007) while, Atta-Allah (1996) and Sharaan *et al* (2002 a) found that the S.C.10 surpassed T.W.C.310 in plant height and leaf area traits.

Table 1. Average values of plant height and leaf area/plant as affected by genotypes, irrigation intervals, nitrogen fertilization and their interactions (Data are combined across seasons)

Treatments	Irrigations	Plant height (cm)				Leaf area / plant (cm ²)			
		Nitrogen				Nitrogen			
Genotypes		N ₁	N ₂	N ₃	Mean	N ₁	N ₂	N ₃	Mean
G ₁	I ₁	212	261	273	248	2320	2377	2436	2378
	I ₂	214	248	261	241	2347	2423	2495	2422
	I ₃	218	251	267	245	2362	2453	2503	2439
Mean		214	253	267	245	2343	2418	2478	2413
G ₂	I ₁	210	247	261	239	2721	2781	2825	2776
	I ₂	203	240	252	231	2760	2811	2880	2817
	I ₃	205	243	256	235	2758	2851	2926	2845
Mean		206	243	256	235	2746	2814	2877	2813
I ₁		211	254	267	244	2521	2579	2631	2577
I ₂		208	244	256	236	2554	2617	2688	2619
I ₃		211	247	262	240	2560	2652	2715	2642
Mean		210	248	262	240	2545	2616	2678	2613

L.S.D 0.05 for:

Genotypes (G):	6.80	9.67
Irrigations (I):	3.11	10.58
(G) x (I):	n.s	n.s
Nitrogen (N):	3.79	8.25
(G) x (N):	n.s	n.s
(I) x (N):	n.s	14.29
(G)x(I)x (N):	n.s	n.s

Application of I₁ treatment significantly increased plant height compared with I₂ or I₃. This result may be attributed to increased availability soil moisture with short irrigation interval. This result are in accordance with El-Ganayni (2000) and Cakir (2004). While application I₃ significantly increased leaf area compared with I₁ or I₂ treatments. The same trend was detected by Yang *et al* (2009) who reported that corn plants under more sever drought stress exhibited

clear improvement for leaf area trait. Using dose of nitrogen fertilization 140 kg N/fed. (N₃) significantly increased plant height and leaf area per plant traits by 262 cm and 2678 cm², respectively, compared to 100 kg N/fed., and 120 kg N /fed., (N₁ and N₂), respectively. The interaction of (I x N) was found to have a significant affect leaf area/plant **Gheysari et al. (2009)** supported this result.

The performance of number of days from planting to mid tasselling and silking are shown in Table (2). Analysis of data indicated that increasing irrigation intervals from 10 (I₁) to 15 (I₂) or 20 (I₃) days caused a significant increase in number of days to 50 % tasselling by 64.50, 65.31 and 65.54 and 50 % silking by 67.46, 68.80 and 69.23, respectively.

Application of 100 Kg N/fed. (N₁) for the two hybrids gave the lowest averages of both number days to 50 % tasselling and silking compared to N₂ and N₃ rates, respectively, where N₁ caused earliness tasselling and silking traits. While the same two traits studied were not affected by the two hybrids. **Masood et al (2003)** found similar results.

The results presented in Table (2) reveal significant interaction between genotypes and irrigation intervals, where planting S.C.10 and T.W.C.310 under I₁ treatment gave the lowest values of number of days to 50% tasselling and silking (64.29 and 66.96 day), respectively. There was also significant interaction among the three factors, where planting maize S.C.10 under I₁ and N₂ treatments gave the lowest averages of number of days 50 % silking (66.75 days) and T.W.C.310 under I₁ and N₁ treatments gave the lowest values of the same trait (67.75 day).

Table 2. Average values of no. of days to 50 % tasselling and silking as affected by genotypes, irrigation intervals, nitrogen fertilization and their interactions (Data are combined across seasons)

Treatments		No. of days to 50 % tasselling				No. of days to 50 % silking			
Genotypes	Irrigations	Nitrogen				Nitrogen			
		N ₁	N ₂	N ₃	Mean	N ₁	N ₂	N ₃	Mean
G ₁	I ₁	63.37	64.38	65.12	64.29	67.25	66.75	66.87	66.96
	I ₂	65.13	66.00	66.12	65.75	69.00	69.12	69.00	69.04
	I ₃	64.25	65.75	66.62	65.54	68.50	68.88	70.62	69.33
Mean		64.25	65.38	65.95	65.19	68.25	68.25	68.83	68.44
G ₂	I ₁	64.00	64.87	65.25	64.71	67.75	68.00	68.13	67.96
	I ₂	64.62	65.00	65.00	64.87	68.37	68.00	69.30	68.56
	I ₃	64.50	65.87	66.25	65.54	68.75	68.87	69.75	69.12
Mean		64.37	65.25	65.50	65.04	68.29	68.29	69.06	68.55
I ₁		63.69	64.63	65.19	64.50	67.50	67.38	67.50	67.46
I ₂		64.88	65.50	65.56	65.31	68.69	68.56	69.15	68.80
I ₃		64.38	65.81	66.44	65.54	68.63	68.88	70.19	69.23
Mean		64.31	65.31	65.73	65.12	68.27	68.27	68.95	68.50

L.S.D 0.05 for:

Genotypes (G):	n.s	n.s
Irrigations (I):	0.412	0.335
(G) x (I):	n.s	0.474
Nitrogen (N):	0.380	0.338
(G) x (N):	n.s	n.s
(I) x (N):	n.s	0.585
(G) x (I) x (N):	n.s	0.828

2-Yield components

Results presented in Table (3) show that ear length and diameter of maize were significantly affected by genotypes, irrigation intervals, nitrogen fertilization. Analysis of variance indicated that genotypes differed significantly in ear length and diameter, where G₁ (S.C.10) had the widest ear diameter and tallest ear length compared to G₂ (T.W.C.310) which may be due to the genetically differences between genotypes. Similar results were obtained by **Sharaan *et al* (2002 a)**.

I₁ application gave the highest values of ear length and diameter i.e. 21.66 and 4.30 cm, respectively. Planting maize with N₁ significantly decreased ear length and diameter by 20.01 and 3.93, respectively, compared with N₂ and N₃. Results in Tale (3) show that planting maize S.C.10 and T.W.C.310 under I₁ gave the longest ear (22.06 and 21.27) and widest diameter (4.45 and 4.14) values, respectively. Whereas, the lowest values were 21.59 and 20.52 (for ear length) 4.24 and 4.01 (for ear diameter), respectively resulted from planting G₁ and G₂ under I₂ treatment.

Table 3. Average values of ear diameter and length as affected by genotypes, irrigation intervals, nitrogen fertilization and their interactions (Data are combined across seasons).

Treatments	Irrigations	Ear diameter (cm)				Ear length (cm)			
		Nitrogen				Nitrogen			
Genotypes		N ₁	N ₂	N ₃	Mean	N ₁	N ₂	N ₃	Mean
G ₁	I ₁	4.17	4.47	4.72	4.45	20.97	22.00	23.20	22.06
	I ₂	3.95	4.27	4.50	4.24	20.46	21.45	22.85	21.59
	I ₃	3.97	4.33	4.55	4.33	20.78	21.76	23.02	21.85
Mean		4.03	4.36	4.59	4.33	20.74	21.74	23.02	21.83
G ₂	I ₁	3.90	4.15	4.37	4.14	20.12	21.37	22.32	21.27
	I ₂	3.80	4.02	4.20	4.01	18.75	20.80	22.00	20.52
	I ₃	3.78	4.06	4.28	4.04	18.97	21.11	22.10	20.73
Mean		3.83	4.08	4.28	4.06	19.28	21.09	22.14	20.84
I ₁		4.04	4.31	4.55	4.30	20.55	21.69	22.76	21.66
I ₂		3.88	4.15	4.35	4.12	19.61	21.13	22.43	21.05
I ₃		3.88	4.20	4.42	4.19	19.88	21.44	22.56	21.29
Mean		3.93	4.22	4.44	4.19	20.01	21.42	22.58	21.34

L.S.D 0.05 for:

Genotypes (G):	0.062	0.132
Irrigations (I):	0.035	0.122
(G) x (I):	0.049	0.173
Nitrogen (N):	0.065	0.217
(G) x (N):	n.s	n.s
(I) x (N):	n.s	0.307
(G) x (I) x (N):	n.s	n.s

Results in Table (4) show that the highest number of rows /ear (12.93) and number of grains /row (43.27) were obtained from planting maize S.C.10 compared with those of T.W.C.310, i.e., 12.51 and 40.14, respectively. The

application of I₁ treatment gave the highest number of rows/ear (12.89) and number of grains/ row (42.40).

Table 4. Average values of number of rows/ear and number of grains/ row as affected by genotypes, irrigation intervals, nitrogen fertilization and their interactions (Data are combined across seasons).

Treatments	Irrigations	Number of rows/ear				Number of grains/ row			
		Nitrogen				Nitrogen			
Genotypes		N ₁	N ₂	N ₃	Mean	N ₁	N ₂	N ₃	Mean
G ₁	I ₁	12.77	13.05	13.55	13.12	42.22	43.35	45.80	43.79
	I ₂	12.20	12.80	13.30	12.77	40.80	42.20	45.07	42.69
	I ₃	12.35	12.92	13.42	12.90	41.76	42.78	45.45	43.33
Mean		12.44	12.92	13.42	12.93	41.59	42.78	45.44	43.27
G ₂	I ₁	12.12	12.60	13.27	12.66	37.17	41.22	44.65	41.01
	I ₂	11.87	12.30	13.12	12.43	35.50	39.77	43.25	39.51
	I ₃	11.97	12.40	13.15	12.51	36.32	40.50	43.60	40.14
Mean		11.99	12.43	13.18	12.53	36.33	40.50	43.83	40.22
I ₁		12.45	12.83	13.41	12.89	39.70	42.29	45.23	42.40
I ₂		12.04	12.55	13.21	12.60	38.15	40.99	44.16	41.10
I ₃		12.16	12.66	13.29	12.70	39.04	41.64	44.53	41.74
Mean		12.21	12.68	13.30	12.73	38.96	41.64	44.64	41.75

L.S.D 0.05 for:

Genotypes (G):	0.081	0.82
Irrigations (I):	0.108	0.26
(G) x (I):	n.s	n.s
Nitrogen (N):	0.140	0.63
(G) x (N):	n.s	0.89
(I) x (N):	n.s	1.09
(G) x (I) x (N):	n.s	n.s

Data given in Table (4) reveal that the N₃ treatment gave the highest number of rows/ear and grains/row traits. The results showed that S.C.10 and T.W.C.310 planted under N₃ treatment produced the highest performance of number of rows 13.42, 13.18 and number of grains 45.44 and 43.83, respectively. Whereas, the genotypes produced the lowest values of the two traits under N₁ treatment.

Data illustrated in Table (5) show that 100-grain weight and grain yield / feddan of S.C.10 outyielded those of T.W.C.310, whereas the values were 35.97 (g) and 2475.37 (kg), respectively. These observations are in full agreement with those of **El- Sheikh (1999) and Sharaan et al (2002 b)**. Results indicated that I₂ caused significant decrease in 100- grain weight and grain yield /feddan, i.e. 34.98 and 2244.93, respectively compared with I₁ treatment. This result may be attributed to the effect of moisture deficit on dry matter accumulation and translocation of metabolites to grains. These results are in harmony with those found by **Mahmood et al (2000), Abo-El- Kheir and Mekki (2007)** who came the same conclusion. The highest values of 100-grain weight and grain

yield/feddan, i.e. 37.53 and 2595.17, respectively, were obtained from N₃ treatment while the lowest values i.e. 32.89 and 1946.90, were obtained from N₁, respectively. In this respect, **Alam *et al* (2003)**, **Masood *et al* (2003)** and **El-Hendawy *et al* (2008)** found that the increasing levels of nitrogen improved the yield and yield components. The interactions of first and second order between the studied variables not attained the level of significance for all cases.

Table 5. Average values of 100-grain weight and grain yield /feddan as affected by genotypes, irrigation intervals, nitrogen fertilization and their interactions (Data are combined across seasons)

Treatments	Irrigations	100-grain weight (g)				Grain yield /feddan(kg)			
		Nitrogen				Nitrogen			
Genotypes		N ₁	N ₂	N ₃	Mean	N ₁	N ₂	N ₃	Mean
G ₁	I ₁	33.80	36.15	38.57	36.17	2116.21	2639.22	2888.18	2547.87
	I ₂	33.28	35.66	38.19	35.71	2041.45	2513.17	2664.36	2406.33
	I ₃	33.75	36.02	38.29	36.02	2053.07	2586.75	2775.96	2471.93
Mean		33.61	35.94	38.35	35.97	2070.24	2579.71	2776.17	2475.37
G ₂	I ₁	32.55	34.75	37.10	34.80	1900.38	2304.21	2467.10	2223.90
	I ₂	31.74	34.59	36.41	34.25	1740.78	2138.50	2371.31	2083.53
	I ₃	32.22	34.41	36.64	34.42	1829.52	2249.16	2404.37	2161.02
Mean		32.17	34.58	36.72	34.49	1823.56	2230.62	2414.26	2156.15
I ₁		33.18	35.45	37.84	35.49	2008.30	2471.72	2677.64	2385.88
I ₂		32.51	35.13	37.30	34.98	1891.12	2325.84	2517.84	2244.93
I ₃		32.99	35.22	37.47	35.22	1941.30	2417.96	2590.17	2316.47
Mean		32.89	35.26	37.53	35.23	1946.90	2405.17	2595.21	2315.76

L.S.D 0.05 for:

Genotypes (G):	0.25	52.76
Irrigations (I):	0.16	20.97
(G) x (I):	n.s	n.s
Nitrogen (N):	0.41	39.14
(G) x (N):	n.s	n.s
(I) x (N):	n.s	n.s
(G) x (I) x (N):	n.s	n.s

REFERENCES

Abo-El-Kheir, M.S.A. and B.B. Mekki (2007). Response of maize single cross-10 to water deficient during silking and grain filling stages. *World J. of Agric. Sci.*3(3):269-272.

Alam, M.M., M.M. Basher, A. Karim, M.A. Rahman and M.R. Ialms (2003). Effect of rate of nitrogen fertilizer and population density on the yield and yield attributes of maize (*Zea mays L.*). *Pakistan J. of Biol. Sci.*6 (20):1770-1773.

- Amanullah, M.J. Hassan, K. Nawab and A. Ali (2007)**. Response of specific leaf area (SLA), leaf area index (LAI) and leaf ratio (LAR) of maize (*Zea mays* L.) to plant density, rate and timing of nitrogen application. *World Appl. Sci. J.* 2 (3):235-243.
- Ashoub, M.A., M.S. Hassanein, I.M.A. Abedel-Azize, M.N. Shahin and M.N. Gohar (1996)**. Influence of irrigation, nitrogen, zinc and manganese fertilization on yield and yield components of maize. *Ann. Agric. Sci., Ain Shams Univ.*, 41(2):697-711.
- Atta-Allah, S.A.A. (1996)**. Effect of irrigation intervals and plant density, growth, yield and its components of some maize varieties. *Proc. Conf. of Agron.*, 9-10 Sept. Cairo:59-70.
- Cakir, R. (2004)**. Effect of water stress at different development stages on vegetative and reproductive growth of corn. *Field Crops Res.* 89:1-16.
- El-Ganayni, A.A. (2000)**. Interaction among irrigation intervals and N levels in corn (*Zea mays* L.). New approach for evaluating interaction. *Mansoura J. Agric. Sci.*, 25(4): 1909-1922.
- El-Hendawy, S.E., E.A. Abd-El-Lattief, M.S. Ahmed and U. Schmidhalter (2008)**. Irrigation rate and plant density effect on yield and water use efficiency of drip-irrigated corn. *Agric. Water Manag.* 95:836-844.
- El-Sheikh, M.H. (1999)**. Response of some maize genotypes to water. *Alex. J. Agric. Res.* 44 (1): 61-77.
- Gheysari, M., S.M. Mirlatifi, M. Bannayan, M. Homae and G. Hoogenboom (2009)**. Interaction of water and nitrogen on maize growth for silage. *Agric. Water Manag.* 96:809-821.
- Gomez, K.A. and A.A. Gomez (1984)**. *Statistical Procedures for Agricultural Research*. 2nd John Wiley and Sons. New York.
- Masood, M., I.H. Shamsi, N. Hussain and W.A. Shah (2003)**. Performance of various maize varieties as affected by different NP levels. *Asian J. plant Sci.* 2 (7):535-538.
- Mohmood, T., M. Saeed and R. Ahmed (2000)**. Impact of water and potassium management on yield and quality of maize (*Zea mays* L.). *Pakistan J. of Biol. Sci.* 3(3): 531-533.
- Sharaan, A.N., F.S. Abd El-Samei, K.M.R. Yousef and H.M. Ibrahim (2002a)**. Response of some maize genotypes to tillage systems and irrigation intervals. 2nd Conf. of Sustain. Agric Develop. 8-10 May, Cairo: 20-30.
- Sharaan, A.N., K.M.R. Yousef, F.S. Abd El-Samei, and H.M. Ibrahim (2002b)**. Maize yield and water relation under combinations of tillage systems and irrigation intervals. 2nd Conf. of Sustain. Agric Develop. 8-10 May, Cairo Univ.,: 31-41.
- Sharar, M.S., M. Ayub, M.A. Nadeem and N. Ahmed (2003)**. Effect of different rates of nitrogen and phosphorus on growth and grain yield of maize (*Zea mays* L.). *Asian J. of Plant Sci.* 2(3):347-349.
- Yang, Y.D.J. Timlin, D.H. Fleisher, S.H. Kim, B. Quebedeaux and V.R. Reddy (2009)**. Simulating leaf area of corn plants at contrasting water status. *Agric. and Forest Meteorology* 149:1161-1167.

إستجابة هجينين من الذرة الشامية لبعض معاملات الإجهاد المانى والتسميد النيتروجينى

فوزى سيد عبد السميع- محمد دسوقى حسن دويدار- سمير كامل على اسماعيل

قسم المحاصيل - كلية الزراعة - جامعة الفيوم - مصر

- أقيمت تجربتان حقلتان بمزرعة كلية الزراعة بالفيوم خلال عامى ٢٠٠٦، ٢٠٠٧ وذلك لتقدير التأثير الفردى والمشارك لثلاث فترات للرى (٢٠، ١٥، ١٠ يوم) و ثلاث معدلات للتسميد النيتروجينى (١٤٠، ١٢٠، ١٠٠ كجم نيتروجين للفدان) على النمو والمحصول ومكوناته لهجينين من الذرة الشامية (هجين فردى ١٠، هجين ثلاثى ٣١٠).
- * أوضحت النتائج تفوق الهجين الفردى ١٠ لمعظم الصفات تحت الدراسة وخصوصاً ارتفاع - وزن المائة حبة ومحصول الحبوب للفدان حيث كانت القيم لهذه الصفات ٢٤٥سم، ٣٥.٩٧جم، ٢٤٧٥.٣٧كجم على الترتيب.
- * أعطت معاملة الرى كل ١٠ أيام زيادة معنوية فى صفات طول النبات - طول الكوز- قطر الكوز- وزن المائة حبة ومحصول الحبوب للفدان.
- * أشارت النتائج إلى أن تطبيق المعدل ١٤٠ كجم نيتروجين للفدان قد أعطى أعلى القيم لصفات طول النبات- مساحة أوراق النبات- عدد الصفوف للكوز- عدد الحبوب فى الصف - وزن المائة حبة والمحصول من الحبوب للفدان بالمقارنة بالمعدلين الأخرين وكانت القيم (٢٦٢سم، ٢٦٧٨سم، ١٣.٣٠ صف ٤٤.٦٤، حبة، ٣٧.٥٣ جم، ٢٥٩٥.١٧ كجم/الفدان) على التوالى.
- * أظهرت النتائج أيضاً أن زراعة الهجين الفردى ١٠ والثلاثى ٣١٠ مع الرى كل ١٠ أيام أعطى أقل القيم فى عدد الايام لظهور ٥٠% من النورات المذكرة و ٥٠% من النورات المؤنثة فى حين كانت زراعة نفس الهجن مع الرى كل ١٠ أيام والمعدل ١٢٠ كجم نيتروجين أظهرت أقل القيم فى عدد الايام لظهور ٥٠% من الأزهار المذكرة.
- * يتضح من إجمالى النتائج أن التأثير الفردى للعوامل تحت الدراسة كان واضحاً على الصفات المدروسة أكثر من التأثير المشترك (التفاعل) الذى لم يصل إلى حد المعنوية فى معظم الصفات - كذلك يجب أن يؤخذ فى الاعتبار دراسة المستويات العالية من التسميد النيتروجينى مع الرى كل ١٠ أيام مع الهجين الفردى ١٠ حيث استجاب هذا الهجين للمستوى العالى ١٤٠ كجم نيتروجين للفدان وأعطى نتائج مرضية من خلال هذه الدراسة.